

ACUTE TOXICITY SUMMARY

CHLOROFORM

(trichloromethane, formyl trichloride, methenyl trichloride, methyl trichloride)

CAS Registry Number: 67-66-3

I. Acute Toxicity Summary (for a 7-hour exposure)

<i>Inhalation reference exposure level</i>	150 µg/m³
<i>Critical effect(s)</i>	histological changes in the nasal epithelium
<i>Hazard Index target(s)</i>	Respiratory System; Nervous System; Reproductive/developmental

II. Physical and Chemical Properties (HSDB, 1993 except as noted)

<i>Description</i>	colorless liquid
<i>Molecular formula</i>	CHCl ₃
<i>Molecular weight</i>	119.49
<i>Density</i>	1.483 g/cm ³ @ 20°C
<i>Boiling point</i>	61°C
<i>Melting point</i>	-63.5°C
<i>Vapor pressure</i>	200 mm Hg @ 25°C
<i>Flashpoint</i>	not applicable; non-flammable liquid, vapor will burn at high temperatures
<i>Explosive limits</i>	not applicable
<i>Solubility</i>	soluble in water, carbon tetrachloride, carbon disulfide, alcohols, benzene, ethers, oils
<i>Odor threshold</i>	192 ppm (geometric mean) (AIHA, 1989)
<i>Odor description</i>	sweet, suffocating (AIHA, 1989)
<i>Metabolites</i>	carbon dioxide, phosgene
<i>Conversion factor</i>	1 ppm = 4.88 mg/m ³ @ 25°C

III. Major Uses or Sources

Chloroform (CHCl₃) is used in industry and laboratory settings as a solvent for adhesives, pesticides, fats, oils, and rubbers. It is also used as a chemical intermediate for fluorocarbon 22, dyes, pesticides, and tribromomethane. It is produced as a byproduct of water and sewage chlorination. Chloroform is also produced in large quantities as a byproduct of wood pulp chlorination in the production of paper products.

IV. Acute Toxicity to Humans

In humans, pulmonary excretion was found to be the major means of elimination following a single oral dose of 0.5 or 1.0 g CHCl_3 (Fry *et al.*, 1972). Up to 68% of the unchanged CHCl_3 and up to 50.6% of the metabolite carbon dioxide were found in the expired air within eight hours of administration. Chloroform in the urine accounted for less than 1% of the oral dose.

Signs of acute CHCl_3 toxicity include fainting, vomiting, dizziness, salivation, fatigue, headache, respiratory depression, and coma (IRIS, 1993). Few reports were found in the literature on the acute toxicity of CHCl_3 to humans in chamber studies. However, a number of case reports exist stemming from its use as an anesthetic.

Cardiac arrhythmia, brachycardia, and cardiac arrest resulting in death have been reported following the use of CHCl_3 as an anesthetic in concentrations of approximately 8,000 to 22,500 ppm (39,000 to 110,000 mg/m^3) (Payne, 1981). Severe liver and kidney damage were noted in an adult male following fatal suicidal ingestion of approximately 6 ounces of CHCl_3 (Piersol *et al.*, 1933).

The incidence of liver enlargement and jaundice was increased in workers exposed to 2-204 ppm (10-995 mg/m^3) CHCl_3 for at least one year (Bomski *et al.*, 1967). Jaundice was reported in 31 workers occupationally exposed to 14-400 ppm (68-1,952 mg/m^3) CHCl_3 for 6 months or less (Phoon *et al.*, 1983).

Predisposing Conditions for Chloroform Toxicity

Medical: Persons with skin, eye, respiratory, liver, kidney or neurological conditions may be more sensitive to the effects of chloroform (Reprotext, 1999).

Chemical: Epinephrine (e.g., in bronchodilators) may potentiate the cardiac effects of chloroform exposure (Reprotext, 1999). Concurrent exposure to barbiturates has been shown to increase chloroform toxicity by induction of liver cytochrome P-450 activity (Cornish *et al.*, 1973). The potentiation of chloroform-induced hepatotoxicity and nephrotoxicity by various alcohols and ketones is well documented (Cowlen *et al.*, 1984; Iijima *et al.*, 1983; Brown and Hewitt, 1984.)

V. Acute Toxicity to Laboratory Animals

Beagle dogs exposed to 14,500 ppm (70,800 mg/m^3) CHCl_3 survived an average of 202 minutes (Von Oettingen *et al.*, 1949). The oral LD_{50} in male and female adult Sprague-Dawley rats is reported as 908 $\text{mg CHCl}_3/\text{kg}$ and 1,117 $\text{mg CHCl}_3/\text{kg}$, respectively (Chu *et al.*, 1980).

Hepatocellular necrosis was observed in adult female mice following a single 4-hour exposure to 200 ppm (976 mg/m^3) CHCl_3 (Kylin *et al.*, 1963). Hepatic fatty infiltration was noted following a single 4-hour exposure to 100 ppm (488 mg/m^3) CHCl_3 . Some studies report that chloroform

Determination of Acute Reference Exposure Levels for Airborne Toxicants
March 1999

renal toxicity is gender-dependent, while hepatotoxicity is similar in both sexes (Smith *et al.*, 1983 and 1984; Hill *et al.*, 1975; Pohl *et al.*, 1984; Taylor *et al.*, 1974).

Cytochrome P-450-mediated metabolism of CHCl_3 in the liver and kidneys has been demonstrated to produce phosgene in rats (Pohl *et al.*, 1979). Hepatotoxicity following chloroform exposure is thought to be due largely to phosgene and other reactive CHCl_3 metabolites. Metabolism of CHCl_3 to phosgene is also responsible for the nephrotoxicity of CHCl_3 (Bailie *et al.*, 1984).

Male rats were exposed to 1, 3, 10, 30, 100, or 300 ppm CHCl_3 6 hours per day for 7 days (Mery *et al.*, 1994). Statistically significant, concentration-dependent, bony proliferation was observed in the ethmoid turbinates of rats exposed to 10 ppm CHCl_3 or greater. Cellular hypertrophy and proliferation in the nasal pharyngeal and olfactory mucosal regions were also increased in a concentration dependent manner in rats exposed to 10 ppm CHCl_3 or greater. No adverse effects were observed following exposure to 3 ppm (15 mg/m^3) CHCl_3 .

VI. Reproductive or Developmental Toxicity

Pregnant rats were exposed to 30, 100, or 300 ppm ($150, 500, \text{ or } 1,500 \text{ mg/m}^3$) CHCl_3 for 7 hours per day on days 6-15 of gestation (Schwetz *et al.*, 1974). A significant increase in the number of fetal resorptions and a decrease in fetal body weights and crown-rump lengths were observed in those animals exposed to 300 ppm CHCl_3 . Following maternal exposure to 100 ppm CHCl_3 , fetuses exhibited a significant increase in malformations including acaudia, imperforate anus, missing ribs and delayed sternal ossification. An increase in the incidence of wavy ribs and delayed skull ossification, as well as reduced fetal crown-rump length, were observed following maternal exposure to 30 ppm CHCl_3 . Maternal toxicity was observed in all three exposure groups.

The incidence of abnormal sperm was significantly increased in male mice exposed to 400 ppm ($1,952 \text{ mg/m}^3$) CHCl_3 for 4 hours/day for 5 days (Land *et al.*, 1981).

Chloroform has not been listed as a developmental or reproductive toxicant under Proposition 65.

VII. Derivation of Acute Reference Exposure Level and Other Severity Levels

Reference Exposure Level (level protective against severe adverse effects; estimated for 7 hour exposure): **0.03 ppm (150 µg/m³)**

<i>Study</i>	Schwetz et al (1974)
<i>Study population</i>	pregnant rats
<i>Exposure method</i>	inhalation exposures to 30, 100, 300 ppm for 7 h/d, days 6-15 of gestation
<i>Critical Effect</i>	fetotoxicity
<i>LOAEL</i>	30 ppm
<i>NOAEL</i>	not determined
<i>Exposure duration</i>	7 hours/day
<i>LOAEL uncertainty factor</i>	10
<i>Interspecies uncertainty factor</i>	10
<i>Intraspecies uncertainty factor</i>	10
<i>Cumulative uncertainty factor</i>	1000
<i>Reference Exposure Level (7 h)</i>	0.03 ppm (0.15 mg/m ³ ; 150 µg/m ³)

The study by Schwetz *et al.* (1974) is the only published developmental toxicity study of chloroform. Exposure of pregnant rats to 30 ppm (150 mg/m³) CHCl₃ for 7 hours per day on days 6-15 of gestation resulted in fetotoxicity as indicated by decreased crown-rump length and increased incidences of wavy ribs and skeletal ossification defects. Maternal toxicity was also observed. An abstract by Dilley *et al.* (1977) indicates an absence of teratological effects in rats exposed to 20,000 mg/m³ CHCl₃ on days 7-14 of gestation. The data from this study were not available for review, therefore, the Schwetz *et al.* study is used in developing the severe adverse effect level for chloroform. A NOAEL was estimated from the reported LOAEL using an uncertainty factor of 10. An additional uncertainty factor of 100 was applied to account for inter- and intraspecies differences. The level protective against severe adverse effects for a 7 hour exposure is estimated as 0.03 ppm (0.15 mg/m³).

Level Protective Against Life-threatening Effects

No recommendation is made due to the limitations of the database. NIOSH (1995) lists a (revised) IDLH of 500 ppm based on acute inhalation toxicity in humans but the selection of the level is somewhat arbitrary and the IDLH does not make allowance for sensitive individuals.

VIII. References

(ATSDR) Agency for Toxic Substances and Disease Registry. Toxicological profile for chloroform. Prepared by Syracuse Research Corporation under subcontract to Clement International Corporation Contract No 205-88-0608. Prepared for US Department of Health and Human Services, Public Health Service, ATSDR; 1991.

Determination of Acute Reference Exposure Levels for Airborne Toxicants
March 1999

(AIHA) American Industrial Hygiene Association. Odor thresholds for chemicals with established occupational health standards. Akron (OH): AIHA; 1989. p. 15.

Bailie MB, Smith JH, Newton JF, Hook JB. Mechanism of chloroform nephrotoxicity. IV. Phenobarbital potentiation of in vitro chloroform metabolism and toxicity in rabbit kidneys. Toxicol Appl Pharmacol 1984;74:285-292.

Bomski H, Sobolewska A, Strakowski A. Toxische Schädigung der Leber durch Chloroform bei Chemiebetriebswerkern [Toxic damage of the liver by chloroform in chemical industry workers]. Int Arch Arbeitsmed 1967;24:127-134.

Brown ES, Hewitt WR. Dose response relationships in ketone-induced potentiation of chloroform hepato- and nephrotoxicity. Toxicol Appl Pharmacol 1984;76:437-453.

Cowlen MS, Hewitt WR, Schroeder F. Mechanisms in 2-hexanone potentiation of chloroform hepatotoxicity. Toxicol Lett 1984;22:293-299.

Chu I, Secours V, Marino L, Villeneuve D. The acute toxicity of four trihalomethanes in male and female rats. Toxicol Appl Pharmacol 1980;52(2):351-353.

Cornish HH, Ling B, Barth M. Phenobarbital and organic solvent toxicity. Am Ind Hyg Assoc J 1973;34:487-492.

Dilley JV, Chernoff N, Kay D, Winslow N, Newell GW. Inhalation teratology studies of five chemicals in rats [abstract]. Toxicol Appl Pharmacol 1977;41:196.

Fry BJ, Taylor T, Hathaway DE. Pulmonary elimination of chloroform and its metabolite in man. Arch Int Pharmacodyn Ther 1972;196:98-111.

Hill RN, Clemens T, Liu D, Vesell E. Genetic control of chloroform toxicity in mice. Science 1975;198:159-160.

(HSDB) Hazardous Substances Data Bank. National Library of Medicine, Bethesda (MD) (CD-ROM Version). Denver (CO): Micromedex, Inc. 1993. (Edition expires 11/31/93).

Iijima M, Cote MG, Plaa GI. A semiquantitative morphologic assessment of chlordecone-potentiated chloroform hepatotoxicity. Toxicol Lett 1983;17:307-314.

(IRIS) Integrated Risk Information System. U.S. Environmental Protection Agency, Washington (DC) (CD-ROM Version). Denver (CO): Micromedex, Inc.; 1993. (Edition expires 11/31/93).

Kylin B, Reichard H, Sumegi L, Yllner S. Hepatotoxicity of inhaled trichloroethylene, tetrachloroethylene, and chloroform: single exposure. Acta Pharmacol Toxicol 1963;20:16-26.

Determination of Acute Reference Exposure Levels for Airborne Toxicants
March 1999

Land PC, Owen E, Linde H. Morphological changes in mouse spermatozoa after exposure to inhalational anesthetics during early spermatogenesis. *Anesthesiology* 1981;54:53-56.

Mery S, Larson JL, Butterworth BE, Wolf DC, Harden R, Morgan KT. Nasal toxicity of chloroform in male F-344 rats and female B6C3F₁ mice following a 1-week inhalation exposure. *Toxicol Appl Pharmacol* 1994;125:214-227.

(NIOSH) National Institute for Occupational Safety and Health. Chemical listing and documentation of revised IDLH values. 1995. (<http://www.cdc.gov/niosh/intridl4.html>)

Payne JP. Chloroform in clinical anaesthesia. *Br J Anaesth* 1981;53 Suppl 1:11s-15s. [cited in ATSDR, 1991.]

Phoon WH, Goh K, Lee L, Tan Kwok S. Toxic jaundice from occupational exposure to chloroform. *Med J Malaysia* 1983;38(1):31-34.

Piersol GM, Tuman H, Kau L. Fatal poisoning following the ingestion of chloroform. *Med Clin N Amer* 1933;17:587-601.

Pohl LR, George JW, Martin JL, Krisha G. Deuterium isotope effect in in vivo bioactivation of chloroform to phosgene. *Biochem Pharmacol* 1979;28:561-563.

Pohl LR, George JW, Satoh H. Strain and sex differences in chloroform-induced nephrotoxicity. Different rates of metabolism of chloroform to phosgene by the mouse kidney. *Drug Metab Dispos* 1984;12(3):304-308.

Reprotext[®] System. Dabney BJ, editor. Denver (CO): Micromedex, Inc.; 1999. (Edition expires 1/31/1999).

Schwetz B, Leong B, Gehring P. Embryo- and fetotoxicity of inhaled chloroform in rats. *Toxicol Appl Pharmacol* 1974;28:442-451.

Smith JH, Maita K, Sleight S, Hook J. Mechanism of chloroform toxicity. 1. Time course of chloroform toxicity in male and female mice. *Toxicol Appl Pharmacol* 1983;70:467-479.

Smith JH, Maita K, Sleight S, Hook J. Effect of sex hormone status on chloroform nephrotoxicity and renal mixed function oxidases in mice. *Toxicology* 1984;30:305-316.

Taylor DC, Brown D, Keeble R, Langley P. Metabolism of chloroform-11. A sex difference in the metabolism of [¹⁴C] chloroform in mice. *Xenobiotica* 1974;4(3):165-174.

Von Oettingen W, Powell C, Alford W, Pecora L. 11 Animal experiments. *NIH Bulletin* #191. 1949;5-6,28-35.